



## PRE-OP PROPHYLAXIS ORDER FORM

All antibiotic administration must be initiated within sixty minutes prior to initial surgical incision or the beginning of the procedure (e.g., introduction of endoscope, insertion of needle, inflation of tourniquet (one hundred twenty (120) minutes prior to incision for Vancomycin.)

Patient Name \_\_\_\_\_ Weight \_\_\_\_\_ kg.

Allergies & Sensitivities \_\_\_\_\_

### Antibiotic Not Applicable

#### Adult Dosing

- CEFAZolin (Ancef) 1 gram IV pre-op (less than or equal to 80 kg)
- CEFAZolin (Ancef) 2 grams IV re-op (greater than 80 kg)
- Clindamycin (Cleocin) 600 mg IV pre-op (infusion over 30 minutes)
- MetroNIDAZOLE (Flagyl) 500 mg IV pre-op (infusion over 20 minutes)
- Gentamicin 80 mg (or \_\_\_\_\_ 1.5 mg/kg) IV pre-op (infusion over 30 minutes)
- Vancomycin 1 gram in 250 ml NSS IV pre-op (less than or equal to 100 kg) (infusion over 60 minutes)  
Reason: \_\_\_\_\_
- Vancomycin 1.5 gram in 500 ml NSS IV pre-op (greater than 100 kg) (infusion over 90 minutes)  
Reason: \_\_\_\_\_

#### Alternative Antibiotic or Pediatric Dosing:

Other Antibiotic: \_\_\_\_\_

Reason: \_\_\_\_\_

#### Additional Orders:

- Sequential Compression Boots       Antiembolic Stockings
- Other: \_\_\_\_\_

Physician Signature \_\_\_\_\_

Date \_\_\_\_\_ Time \_\_\_\_\_

Acknowledged by RN Signature \_\_\_\_\_

Date \_\_\_\_\_ Time \_\_\_\_\_